

hep-ph/0306147
KEK-TH-894
February 1, 2008

A Possibility of large electro-weak penguin contribution in $B \rightarrow K\pi$ modes

Tadashi Yoshikawa

Theory Group, KEK Tsukuba, 305-0801, Japan.

Abstract

We discuss about a possibility of large electro-weak penguin contribution in $B \rightarrow K\pi$ from recent experimental data. The several relations among the branching ratios which realize when the contributions from tree type and electro-weak penguin are small compared with the gluon penguin and can be treated as the expansion parameters do not satisfy the data. The difference comes from the r^2 terms which is the square of the ratio with the gluon penguin and the main contribution comes from electro-weak penguin. We find that the contribution from electro-weak penguin may be large to explain the experimental data. If the magnitude estimated from experiment is quite large compared with the theoretical estimation, then it may be including some new physics effects.

One of the main targets at the B factories is to determine the all CP angles in the unitarity triangles of the Cabbibo-Kobayashi-Maskawa (CKM) matrix [1]. ϕ_1 [2] as one of the angles has already been measured and established the CP violation in the B meson system by Belle[3] and BaBar[4] collaborations. The next step is to measure the remaining angles. The canonical decay modes for measuring ϕ_2 and ϕ_3 are $B_d^0 \rightarrow \pi^+\pi^-$ and $B^\pm \rightarrow DK^\pm$ respectively but the methods have some difficulties to extract cleanly the angles. To avoid the difficulty, isospin relation[6] and $SU(3)$ relation including $B \rightarrow K\pi$ modes [7]-[10] are being considered as a method to extract the weak phases.

$B \rightarrow K\pi$ modes has been also measured[5] and they will be useful informations to understand the CP violation through the KM phases. If we can directly solve about these modes, it is very elegant way to determine the parameters and the weak phase. However we can not do so because there are too many parameters in $B \rightarrow K\pi$ modes to extract the weak phases. So we need understand these modes step by step. To understand the weak phase through this mode, there are several approaches by diagram decomposition[7]-[13], QCD factorization[14] and pQCD[15] and so on. The contributions including the weak phase ϕ_3 come from tree type diagrams which has CKM suppression factor and they are usually dealt with small parameter compared with gluon penguin. If we can deal the contributions except for gluon penguin with the small parameters, then, there are several relations among the averaged branching ratios of $B \rightarrow K\pi$ modes. For example, $Br(K^+\pi^-)/2Br(K^0\pi^0) \approx 2Br(K^+\pi^0)/Br(K^0\pi^+)$ [14] et al. However, the recent experiment does not seem to satisfy them so well. When we reconsider these modes to compare with the data, we find that the role of a color favored electro-weak penguin may be important to explain the difference between the relations and the experimental data. The color favored electro-weak penguin diagram is included in $B^0 \rightarrow K^0\pi^0$ and $B^0 \rightarrow K^+\pi^0$ and the data of the branching ratio are slightly larger than half of $B^0 \rightarrow K^+\pi^-$, where the 1/2 comes from the difference of π^0 and π^+ in final state. So we need to know the informations about the electro-weak penguin contributions in $B \rightarrow K\pi$ decay modes to understand about the effect from the weak phases. The role was pointed out and the magnitude was estimated in several works[14]. They said that the ratio between gluon and electro-weak penguins is about 0.14 as the central value but the experimental data may suggest that the magnitude seem to be slightly larger than the estimation. If there is quite large deviation in the contribution from the electro-weak penguin, it may suggest a possibility of new physics in these modes.

In $B^0 \rightarrow K_S\pi^0$, we can consider to use the time-dependent CP asymmetry to extract the weak phase. However the mode has the electro-weak penguin diagram so that one must remove the contribution to extract ϕ_3 . And we have to check whether extracting ϕ_3 is possible or not. To do so, it is important to reconsider about the electro-weak penguin.

In this work, we consider how large contribution from the electro-weak penguins from only experimental data.

The decay amplitudes of $B \rightarrow K\pi$ are

$$\begin{aligned} A^{0+} &\equiv A(B^+ \rightarrow K^0\pi^+) \\ &= \left[AV_{ub}^*V_{us} + \sum_{i=u,c,t} (P_i + EP_i - \frac{1}{3}P_{EWi} + \frac{2}{3}EP_{EWi}^C)V_{ib}^*V_{is} \right], \end{aligned} \quad (1)$$

$$\begin{aligned}
A^{00} &\equiv A(B^0 \rightarrow K^0 \pi^0) \\
&= -\frac{1}{\sqrt{2}} \left[CV_{ub}^* V_{us} - \sum_{i=u,c,t} (P_i + EP_i - P_{EWi} - \frac{1}{3}P_{EWi}^C - \frac{1}{3}EP_{EWi}^C) V_{ib}^* V_{is} \right], \quad (2)
\end{aligned}$$

$$\begin{aligned}
A^{+-} &\equiv A(B^0 \rightarrow K^+ \pi^-) \\
&= -\left[TV_{ub}^* V_{us} + \sum_{i=u,c,t} (P_i + EP_i + \frac{2}{3}P_{EWi}^C - \frac{1}{3}EP_{EWi}^C) V_{ib}^* V_{is} \right], \quad (3)
\end{aligned}$$

$$\begin{aligned}
A^{+0} &\equiv A(B^+ \rightarrow K^+ \pi^0) \\
&= -\frac{1}{\sqrt{2}} [(T + C + A) V_{ub}^* V_{us} \\
&\quad + \sum_{i=u,c,t} (P_i + EP_i + P_{EWi} + \frac{2}{3}P_{EWi}^C + \frac{2}{3}EP_{EWi}^C) V_{ib}^* V_{is}] , \quad (4)
\end{aligned}$$

where T is a color favored tree amplitude, C is a Color suppressed tree, A is an annihilation, $P_i (i = u, c, t)$ is a gluonic penguin, EP_i is a penguin exchange, P_{EWi} is a color favored electroweak penguin, P_{EWi}^C is a color suppressed electroweak penguin and EP_{EWi}^C is a color suppressed electro-weak penguin exchange diagram. After this, for simplicity, we neglect the u - and c - electroweak penguin diagram because of the smallness, And we redefine the each terms as following

$$T + P_u + EP_u - P_c - EP_c \rightarrow T, \quad (5)$$

$$C - P_u - EP_u + P_c + EP_c \rightarrow C, \quad (6)$$

$$A + P_u + EP_u - P_c - EP_c \rightarrow A, \quad (7)$$

$$P_t + EP_t - P_c - EP_c - \frac{1}{3}P_{EW}^C + \frac{2}{3}EP_{EW}^C \rightarrow P, \quad (8)$$

$$P_{EW} + EP_{EW}^C \rightarrow P_{EW}, \quad (9)$$

$$P_{EW}^C - EP_{EW}^C \rightarrow P_{EW}^C. \quad (10)$$

One can reduce the number of parameter up to 6 (or 12). Then, the amplitudes are, by using the unitarity relation of KM matrix,

$$A^{0+} = [PV_{tb}^* V_{ts} + AV_{ub}^* V_{us}], \quad (11)$$

$$A^{00} = \frac{1}{\sqrt{2}} [(P - P_{EW}) V_{tb}^* V_{ts} - CV_{ub}^* V_{us}], \quad (12)$$

$$A^{+-} = -[(P + P_{EW}^C) V_{tb}^* V_{ts} + TV_{ub}^* V_{us}], \quad (13)$$

$$A^{+0} = -\frac{1}{\sqrt{2}} [(P + P_{EW} + P_{EW}^C) V_{tb}^* V_{ts} + (T + C + A) V_{ub}^* V_{us}]. \quad (14)$$

By this diagram decomposition[8], one can easily find the isospin relation among the amplitudes,

$$\sqrt{2}A^{+0} + A^{0+} = \sqrt{2}A^{00} + A^{+-}. \quad (15)$$

They are rewritten as follows:

$$A^{0+} = -P|V_{tb}^*V_{ts}| \left[1 - r_A e^{i\delta^A} e^{i\phi_3} \right], \quad (16)$$

$$A^{00} = -\frac{1}{\sqrt{2}}P|V_{tb}^*V_{ts}| \left[1 - r_{EW} e^{i\delta^{EW}} + r_C e^{i\delta^C} e^{i\phi_3} \right], \quad (17)$$

$$A^{+-} = P|V_{tb}^*V_{ts}| \left[1 + r_{EW}^C e^{i\delta^{EW^C}} - r_T e^{i\delta^T} e^{i\phi_3} \right], \quad (18)$$

$$A^{+0} = \frac{1}{\sqrt{2}}P|V_{tb}^*V_{ts}| \left[1 + r_{EW} e^{i\delta^{EW}} + r_{EW}^C e^{i\delta^{EW^C}} - (r_T e^{i\delta^T} + r_C e^{i\delta^C} + r_A e^{i\delta^A}) e^{i\phi_3} \right], \quad (19)$$

where ϕ_3 is the weak phase of $V_{ub}^*V_{us}$, δ^X is the strong phase difference between each diagram and gluon penguin and

$$r_A = \frac{|AV_{ub}^*V_{us}|}{|PV_{tb}^*V_{ts}|}, \quad r_T = \frac{|TV_{ub}^*V_{us}|}{|PV_{tb}^*V_{ts}|}, \quad r_C = \frac{|CV_{ub}^*V_{us}|}{|PV_{tb}^*V_{ts}|}, \quad (20)$$

$$r_{EW} = \frac{|P_{EW}|}{|P|}, \quad r_{EW}^C = \frac{|P_{EW}^C|}{|P|}. \quad (21)$$

We assume as the hierarchy of the ratios that $1 > r_T, r_{EW} > r_C, r_{EW}^C > r_A$ [8]. $|P/T|$ was estimated about 0.1 in [19]¹ by considering the $B \rightarrow \pi\pi$ and it was also shown by the ratio of branchings of $B^+ \rightarrow \pi^0\pi^+$ and $B^+ \rightarrow K^0\pi^+$ [16, 17, 18]. In $B \rightarrow K\pi$ mode, the tree type diagram is suppressed by KM factor $V_{ub}^*V_{us}$ and $r_T \sim |T/P| \times \lambda^2 R_b \sim 0.2$, where cabbibo angle $\lambda = 0.22$ and we used $R_b = \sqrt{\rho^2 + \eta^2} \sim 0.4$. r_C and r_{EW}^C are suppressed by color factor from r_T and r_{EW} . Comparing the Wilson coefficients which correspond to the diagrams under the factorization method, we assume that $r_C \sim 0.1r_T$ and $r_{EW}^C \sim 0.1r_{EW}$ [16, 14]. Here we do not put any assumption about the magnitude of r_{EW} . r_A could be negligible because it should have B meson decay constant and it works as a suppression factor f_B/M_B . According to this assumption, we will neglect the r^2 terms including r_C, r_A and r_{EW}^C . Then, the averaged branching ratios are

$$\begin{aligned} \bar{B}^{0+} &\propto \frac{1}{2} \left[|A^{0+}|^2 + |A^{0-}|^2 \right] \\ &= |P|^2 |V_{tb}^*V_{ts}|^2 \left[1 - 2r_A \cos \delta^A \cos \phi_3 \right], \end{aligned} \quad (22)$$

$$\begin{aligned} 2\bar{B}^{00} &\propto \left[|A^{00}|^2 + |\bar{A}^{00}|^2 \right] \\ &= |P|^2 |V_{tb}^*V_{ts}|^2 \left[1 + r_{EW}^2 - 2r_{EW} \cos \delta^{EW} + 2r_C \cos \delta^C \cos \phi_3 \right], \end{aligned} \quad (23)$$

$$\begin{aligned} \bar{B}^{+-} &\propto \frac{1}{2} \left[|A^{+-}|^2 + |A^{-+}|^2 \right] \\ &= |P|^2 |V_{tb}^*V_{ts}|^2 \left[1 + r_T^2 + 2r_{EW}^C \cos \delta^{EW^C} - 2r_T \cos \delta^T \cos \phi_3 \right], \end{aligned} \quad (24)$$

$$2\bar{B}^{+0} \propto \left[|A^{+0}|^2 + |A^{-0}|^2 \right]$$

¹Note that this ratio $|P/T|$ does not include CKM factors.

	CLEO[21]	Belle[5, 22]	BaBar[23, 24]	Average
$Br(B^+ \rightarrow K^0 \pi^+) \times 10^6$	$18.8^{+3.7+2.1}_{-3.3-1.8}$	$22.0 \pm 1.9 \pm 1.1$	$17.5^{+1.8}_{-1.7} \pm 1.3$	19.6 ± 1.4
$Br(B^0 \rightarrow K^0 \pi^0) \times 10^6$	$12.8^{+4.0+1.7}_{-3.3-1.4}$	$12.6 \pm 2.4 \pm 1.4$	$10.4 \pm 1.5 \pm 0.8$	11.2 ± 1.4
$Br(B^0 \rightarrow K^+ \pi^-) \times 10^6$	$18.0^{+2.3+1.2}_{-2.1-0.9}$	$18.5 \pm 1.0 \pm 0.7$	$17.9 \pm 0.9 \pm 0.7$	18.2 ± 0.8
$Br(B^+ \rightarrow K^+ \pi^0) \times 10^6$	$12.9^{+2.4+1.2}_{-2.2-1.1}$	$12.8 \pm 1.4^{+1.4}_{-1.0}$	$12.8^{+1.2}_{-1.1} \pm 1.0$	12.8 ± 1.1

Table 1: The experimental data and the average.

$$\begin{aligned}
= & |P|^2 |V_{tb}^* V_{ts}|^2 \left[1 + r_{EW}^2 + r_T^2 + 2r_{EW} \cos \delta^{EW} + 2r_{EW}^C \cos \delta^{EWC} \right. \\
& - (2r_T \cos \delta^T + 2r_C \cos \delta^C + 2r_A \cos \delta^A) \cos \phi_3 \\
& \left. - 2r_{EW} r_T \cos(\delta^{EW} - \delta^T) \cos \phi_3 \right]. \quad (25)
\end{aligned}$$

One can take several ratios between the branching ratios. If all modes are gluon penguin dominant, the ratios should be close to 1. The shift from 1 will depend on the magnitude of r s. From the averaged values of the recent experimental data in Table 1,

$$\frac{\bar{B}^{+-}}{2\bar{B}^{00}} = 0.81 \pm 0.11, \quad (26)$$

$$\frac{2\bar{B}^{+0}}{\bar{B}^{0+}} = 1.31 \pm 0.15, \quad (27)$$

$$\frac{\tau^+ \bar{B}^{+-}}{\tau^0 \bar{B}^{0+}} = 1.01 \pm 0.09, \quad (28)$$

$$\frac{\tau^0 \bar{B}^{+0}}{\tau^+ \bar{B}^{00}} = 1.05 \pm 0.16, \quad (29)$$

$$\frac{\tau^+ 2\bar{B}^{00}}{\tau^0 \bar{B}^{0+}} = 1.24 \pm 0.18, \quad (30)$$

$$\frac{\tau^0 2\bar{B}^{+0}}{\tau^+ \bar{B}^{+-}} = 1.30 \pm 0.13, \quad (31)$$

where $\frac{\tau^+}{\tau^0}$ is a lifetime ratio between the charged and the neutral B mesons and $\tau(B^\pm)/\tau(B^0) = 1.083 \pm 0.017$ [20]. While, from eq.(22)-(25) under the assumption that all r is smaller than 1 and the r^2 terms including r_C, r_A and r_{EW}^C are neglected,

$$\begin{aligned}
\frac{\bar{B}^{+-}}{2\bar{B}^{00}} = & \left\{ 1 + 2r_{EW} \cos \delta^{EW} + 2r_{EW}^C \cos \delta^{EWC} - 2(r_T \cos \delta^T + r_C \cos \delta^C) \cos \phi_3 + r_T^2 \right\} \\
& - r_{EW}^2 + 4r_{EW}^2 \cos^2 \delta^{EW}, \quad (32)
\end{aligned}$$

$$\begin{aligned}
\frac{2\bar{B}^{+0}}{\bar{B}^{0+}} = & \left\{ 1 + 2r_{EW} \cos \delta^{EW} + 2r_{EW}^C \cos \delta^{EWC} - 2(r_T \cos \delta^T + r_C \cos \delta^C) \cos \phi_3 + r_T^2 \right\} \\
& + r_{EW}^2 - 2r_{EW} r_T \cos(\delta^{EW} - \delta^T) \cos \phi_3, \quad (33)
\end{aligned}$$

$$\frac{\tau^+ \bar{B}^{+-}}{\tau^0 \bar{B}^{0+}} = 1 + 2r_{EW}^C \cos \delta^{EWC} - 2(r_T \cos \delta^T - r_A \cos \delta^A) \cos \phi_3 + r_T^2, \quad (34)$$

$$\begin{aligned} \frac{\tau^0 \bar{B}^{+0}}{\tau^+ \bar{B}^{00}} &= 1 + 2r_{EW}^C \cos \delta^{EWC} - 2(r_T \cos \delta^T + 2r_C \cos \delta^C + r_A \cos \delta^A) \cos \phi_3 + r_T^2 \\ &\quad + 4r_{EW} \cos \delta^{EW} - 2r_{EW} r_T \cos(\delta^{EW} - \delta^T) \cos \phi_3 + 4r_{EW}^2 \cos^2 \delta^{EW}, \end{aligned} \quad (35)$$

$$\frac{\tau^+ 2\bar{B}^{00}}{\tau^0 \bar{B}^{0+}} = 1 - 2r_{EW} \cos \delta^{EW} - 2(r_C \cos \delta^C + r_A \cos \delta^A) \cos \phi_3 + r_{EW}^2. \quad (36)$$

$$\begin{aligned} \frac{\tau^0 2\bar{B}^{+0}}{\tau^+ \bar{B}^{+-}} &= 1 + 2r_{EW} \cos \delta^{EW} - 2(r_C \cos \delta^C + r_A \cos \delta^A) \cos \phi_3 + r_{EW}^2 \\ &\quad - 2r_{EW} r_T \cos(\delta^{EW} - \delta^T) \cos \phi_3 + 4r_T^2 \cos^2 \delta^T \cos^2 \phi_3. \end{aligned} \quad (37)$$

If all modes are dominated by only gluonic penguin contribution, namely, all r is much smaller than 1, then all ratio of branching ratios should be 1. But the data are not so. eqs.(32),(33), (36) and (37) seem to differ from 1 so that there must be some sizable contributions except for the gluon penguin.

If we can neglect all r^2 terms, then there are a few relations among eqs.(32)-(37) as following

$$\frac{\bar{B}^{+-}}{2\bar{B}^{00}} = \frac{2\bar{B}^{+0}}{\bar{B}^{0+}}, \quad (38)$$

$$\frac{2\bar{B}^{+0}}{\bar{B}^{0+}} - \frac{\tau^+ \bar{B}^{+-}}{\tau^0 \bar{B}^{0+}} + \frac{\tau^+ 2\bar{B}^{00}}{\tau^0 \bar{B}^{0+}} - 1 = 0. \quad (39)$$

However, the experimental data listed in eqs.(26)-(31) do not satisfy these relations so well. According to the experimental data, $\frac{\bar{B}^{+-}}{2\bar{B}^{00}}$ seems to be smaller than 1 but $\frac{2\bar{B}^{+0}}{\bar{B}^{0+}}$ be larger than 1. So it shows there is a discrepancy between them. The equations of $\frac{\bar{B}^{+-}}{2\bar{B}^{00}}$ and $\frac{2\bar{B}^{+0}}{\bar{B}^{0+}}$ are same up to r_T^2 term and the difference comes from r^2 term including r_{EW} . r_T^2 term does not seem to contribute to the ratios so strongly. The second relation corresponds to the isospin relation of eq.(15) at the first order of r . The discrepancy of relation (39) from 0 also comes from r^2 term including r_{EW} . The differences are

$$\frac{2\bar{B}^{+0}}{\bar{B}^{0+}} - \frac{\bar{B}^{+-}}{2\bar{B}^{00}} = 2r_{EW}^2 - 2r_{EW} r_T \cos(\delta^{EW} - \delta^T) \cos \phi_3 - 4r_{EW}^2 \cos^2 \delta^{EW} = 0.50 \pm 0.19, \quad (40)$$

$$\frac{2\bar{B}^{+0}}{\bar{B}^{0+}} - \frac{\tau^+ \bar{B}^{+-}}{\tau^0 \bar{B}^{0+}} + \frac{\tau^+ 2\bar{B}^{00}}{\tau^0 \bar{B}^{0+}} - 1 = 2r_{EW}^2 - 2r_{EW} r_T \cos(\delta^{EW} - \delta^T) \cos \phi_3 = 0.54 \pm 0.25, \quad (41)$$

so that one can find the electro-weak penguin contributions may be large. All terms are including r_{EW} and the deviation of the relation from 0 is finite. Here the error are determined by adding quadratically all errors. Using the other relation as following

$$\begin{aligned} \frac{\bar{B}^{+-}}{2\bar{B}^{00}} - \frac{\tau^0 \bar{B}^{+0}}{\tau^+ \bar{B}^{00}} + \frac{\tau^+ 2\bar{B}^{00}}{\tau^0 \bar{B}^{0+}} - 1 \\ = -4r_{EW} \cos \delta^C + 2r_{EW} r_T \cos(\delta^{EW} - \delta^T) \cos \phi_3 = 0.00 \pm 0.26, \end{aligned} \quad (42)$$

we can solve them about r_{EW}^2 and if we can respect the central values, the solutions are

$$(r_{EW}, \cos \delta^{EW}, r_T \cos(\delta^{EW} - \delta^T) \cos \phi_3) = (0.26, -0.38, -0.75) \text{ and } (0.69, 0.21, 0.41). \quad (43)$$

This solution show that large electro-weak penguin contribution (but $r_T \cos(\delta^{EW} - \delta^T) \cos \phi_3$ is too large because r_T was estimated around 0.2 by the other methods.) The allowed region of r_{EW} , $\cos \delta^{EW}$ and $r_T \cos(\delta^{EW} - \delta^T) \cos \phi_3$ at 1σ level for eqs.(40)-(42) is shown in Fig.1.




Figure 1: The allowed region on $(r_{EW}, \cos \delta^{EW})$ and $(r_{EW}, r_T \cos(\delta^{EW} - \delta^T) \cos \phi_3)$ plane at 1σ level.

From this result, we find that the smaller r_{EW} will favored a larger $|r_T \cos(\delta^{EW} - \delta^T) \cos \phi_3|$ term with negative sign. However such large r_T is disfavored by the rough estimation r_T should be around 0.2 which will satisfy that $|P/T| \sim 0.1$ to explain the large CP violation in $B^0 \rightarrow \pi^+ \pi^-$. Even if $|r_T \cos(\delta^{EW} - \delta^T) \cos \phi_3|$ is within 0.2, then r_{EW} will also be larger than 0.2 and r_{EW} will be larger than r_T . This is showing that there is a possibility of large electro-weak penguin contribution. Note that in the case r_{EW} is quite large, the expansion by r_{EW} may not be good but eq.(41) will be satisfied. Roughly speaking, the shift of eqs.(26)-(31) from 1 seem to depend on the r_{EW}^2 term and the sign. To fix the solution or confirm the large electro-weak penguin contribution, we need higher accurate data.

The contributions from tree diagram are not so large except for the cross term with the electro-weak penguin because $\frac{\tau^+ \bar{B}^{+-}}{\tau^0 \bar{B}^{0+}}$ is quite near 1. When we consider the direct CP asymmetry, the experimental data in Table. 2 do not also show so large value. The CP asymmetries under the same assumption are

$$A_{CP}^{0+} \equiv \frac{|A^{0-}|^2 - |A^{0+}|^2}{|A^{0-}|^2 + |A^{0+}|^2} = -2r_A \sin \delta^A \sin \phi_3 \sim 0.0, \quad (44)$$

$$A_{CP}^{00} \equiv \frac{|\bar{A}^{00}|^2 - |A^{00}|^2}{|\bar{A}^{00}|^2 + |A^{00}|^2} = 2r_C \sin \delta^C \sin \phi_3, \quad (45)$$

$$A_{CP}^{+-} \equiv \frac{|A^{-+}|^2 - |A^{+-}|^2}{|A^{-+}|^2 + |A^{+-}|^2} = -2r_T \sin \delta^T \sin \phi_3 - r_T^2 \sin 2\delta^T \sin 2\phi_3, \quad (46)$$

	Belle[5, 25]	BaBar[23, 24]	Average
$A_{CP}(B^+ \rightarrow K^0 \pi^+)$	$0.07^{+0.09+0.01}_{-0.08-0.03}$	$-0.17 \pm 0.10 \pm 0.02$	-0.03 ± 0.07
$A_{CP}(B^0 \rightarrow K^0 \pi^0)$	-	$0.03 \pm 0.036 \pm 0.09$	0.03 ± 0.37
$A_{CP}(B^0 \rightarrow K^+ \pi^-)$	$-0.07 \pm 0.06 \pm 0.01$	$-0.10 \pm 0.05 \pm 0.02$	-0.09 ± 0.04
$A_{CP}(B^+ \rightarrow K^+ \pi^0)$	$0.23 \pm 0.11^{+0.01}_{-0.04}$	$-0.09 \pm 0.09 \pm 0.01$	0.04 ± 0.07

Table 2: The experimental data of the direct CP asymmetry and the average.

$$\begin{aligned}
A_{CP}^{+0} \equiv \frac{|A^{-0}|^2 - |A^{+0}|^2}{|A^{-0}|^2 + |A^{+0}|^2} = & -2(r_T \sin \delta^T + r_C \sin \delta^C + r_A \sin \delta^A) \sin \phi_3 \\
& - 2r_{EW} r_T \sin(\delta^T - \delta^{EW}) \sin \phi_3 - r_T^2 \sin 2\delta^T \sin 2\phi_3 \\
& + 2r_{EW} r_T \sin \delta^T \cos \delta^{EW} \sin \phi_3.
\end{aligned} \tag{47}$$

A_{CP}^{+0} should be almost 0 because of the smallness of r_A and the data is also showing it. Up to the first order of r , there is a relation among the CP asymmetries as follows:

$$A_{CP}^{+0} = A_{CP}^{+-} - A_{CP}^{00} + A_{CP}^{0+}. \tag{48}$$

The discrepancy of this relation is also caused by the cross term of r_T and r_{EW} . If we can have more accurate data, this may also show an useful information about r_{EW} . Because A_{CP}^{+-} is not so large, we can confirm that r_T will not become so large value.

In $B \rightarrow \pi^0 K_S$, we can also use some informations about time dependent CP asymmetry[26]. The measurements for $B \rightarrow \pi^0 K_S$ are

$$\Gamma(B^0 \rightarrow \pi^0 K_S) + \Gamma(\bar{B}^0 \rightarrow \pi^0 K_S) \propto (|A|^2 + |\bar{A}|^2), \tag{49}$$

$$\begin{aligned}
\Gamma(B^0(t) \rightarrow \pi^0 K_S) - \Gamma(\bar{B}^0(t) \rightarrow \pi^0 K_S) \propto & (|A|^2 - |\bar{A}|^2) \cos \Delta mt \\
& + 2Im(e^{-2i\phi_1} A^* \bar{A}) \sin \Delta mt.
\end{aligned} \tag{50}$$

After neglecting the contribution from annihilation diagram, we consider the following measurements under the same assumption,

$$\begin{aligned}
\frac{\tau^+ \Gamma(B^0 \rightarrow \pi^0 K_S) + \Gamma(\bar{B}^0 \rightarrow \pi^0 K_S)}{\tau^0 \Gamma(B^+ \rightarrow \pi^+ K_S)} & \equiv X \\
& = 1 - 2r_{EW} \cos \delta^{EW} + r_{EW}^2 + 2r_C \cos \phi_3 \cos \delta^C,
\end{aligned} \tag{51}$$

$$\begin{aligned}
\frac{\tau^+ \Gamma(B^0(t) \rightarrow \pi^0 K_S) - \Gamma(\bar{B}^0(t) \rightarrow \pi^0 K_S)}{\tau^0 \Gamma(B^+ \rightarrow \pi^+ K_S)} & \equiv Y \cos \Delta mt - Z \sin \Delta mt \\
& = (-2r_C \sin \phi_3 \sin \delta^C) \cos \Delta mt \\
& \quad - \{\sin 2\phi_1 (1 - 2r_{EW} \cos \delta^{EW} + r_{EW}^2) \\
& \quad + 2r_C \sin(\phi_3 + 2\phi_1) \cos \delta^C\} \sin \Delta mt.
\end{aligned} \tag{52}$$

And we define them as follows:

$$X = 1 - 2r_{EW} \cos \delta^{EW} + r_{EW}^2 + 2r_C \cos \phi_3 \cos \delta^C, \tag{53}$$

$$Y = -2r_C \sin \phi_3 \sin \delta^C \approx -A_{CP}^{00}, \quad (54)$$

$$Z = \sin 2\phi_1 (1 - 2r_{EW} \cos \delta^{EW} + r_{EW}^2) + 2r_C \sin (\phi_3 + 2\phi_1) \cos \delta^C. \quad (55)$$

Eliminating $r_C \cos \delta^C$ in X and Z , we find $\tan \phi_3$ as a function of r_{EW} ,

$$\tan \phi_3 = \frac{\{Z - X \sin 2\phi_1\}}{\cos 2\phi_1 \{X - (1 - 2r_{EW} \cos \delta^{EW} + r_{EW}^2)\}}. \quad (56)$$

If one can estimate r_{EW} at good accuracy, one can have a information about ϕ_3 by inputting the value of ϕ_1 which is measured by $B \rightarrow J/\psi K_S$ at higher accurate experiment. It will be extracted from quantity of $O(0.01)$ because $r_C \sim 0.02$ under the assumption in this work. Indeed, for the solution eq.(43), $2r_C \cos \delta^C \cos \phi_3 \approx X - 1 + 2r_{EW} \cos \delta^{EW} - r_{EW}^2 \sim -0.03$. So to use this measurements one may need some corrections from $K - \bar{K}$ mixing and the width difference. See Appendix. After extracting ϕ_3 by the other modes, we can use to estimate or confirm how large is the electro-weak penguin contribution.

In this paper, we discussed about a possibility of large electro-weak penguin contribution in $B \rightarrow K\pi$ from recent experimental data. The several relations among the branching ratios which realize when the contributions from tree type and electro-weak penguin are small compared with the gluon penguin do not satisfy the data. The difference comes from the r^2 terms and the main contribution comes from electro-weak penguin. We find that the contribution from electro-weak penguin may be larger than from tree diagrams to explain the experimental data. If the magnitude estimated from experiment is quite large compared with the theoretical estimation which is usually smaller than tree contributions[12, 13, 14], then it may be including some new physics effects. In this analysis, we find that who can have some contribution from new physics is the color favored electro-weak penguin type diagram which is the process π^0 goes out from $B - K$ current.

Acknowledgments

We would like to thank Sechul Oh for many useful comments and discussions.

Appendix: Some correction terms from the $K - \bar{K}$ mixing and the width difference.

In the discussion about the time dependent CP asymmetry of $B^0 \rightarrow K_S \pi^0$, we neglected the effect of CP violation in K meson system and tiny width difference of B_d mesons because it is very small. The magnitude is $|\epsilon| = 2 \times 10^{-3}$ and $\Delta\Gamma_d/\Gamma_d$ is about 3×10^{-3} which is estimated in [27]. If r_C is the order, we must deal with the contributions. In the Kon system, we define the K_S and K_L mesons as follows:

$$|K^0\rangle = \frac{1-\epsilon}{\sqrt{2}} [|K_S\rangle + |K_L\rangle], \quad (57)$$

$$|\bar{K}^0\rangle = \frac{1+\epsilon}{\sqrt{2}} [|K_S\rangle - |K_L\rangle], \quad (58)$$

where ϵ is the parameter which shows the CP violation of K. Then one needs several correction terms including ϵ and $\Delta\Gamma_d$ as the expansion parameters[27] in X , Y and Z (Note that the definition of the sign of ϵ is different from it in [27].) as follows:

$$X(t) = 1 - 2r_{EW} \cos \delta^{EW} + r_{EW}^2 + 2r_C \cos \phi_3 \cos \delta^C + \cos 2\phi_1 \sinh \frac{\Delta\Gamma_d t}{2} + \dots, \quad (59)$$

$$Y = 2r_C \sin \phi_3 \sin \delta^C - 2Re(\epsilon) + O(\epsilon r_C), \quad (60)$$

$$Z = \sin 2\phi_1 (1 - 2r_{EW} \cos \delta^{EW} + r_{EW}^2) + 2r_C \sin(\phi_3 + 2\phi_1) \cos \delta^C - 2Im(\epsilon) \cos 2\phi_1 + \sin 2\phi_1 \cos 2\phi_1 \sinh \frac{\Delta\Gamma_d t}{2} + O(\epsilon r_C). \quad (61)$$

Here we neglected the terms of ϵr . In addition to these terms, there are also a constant term and a proportional term to $\sin^2(\Delta mt)$. X has also a correction term by ϵ but it is of order ϵr so that one can neglect them. However, indeed, the correction terms has already been included in $\sin 2\phi_1$ determined by $B \rightarrow J/\psi K_S$ [27] and the value which subtracts from Z is effectively $(\sin 2\phi_1 - 2Im(\epsilon) \cos 2\phi_1 + \dots)$ so that one can neglect the effect in Z .

References

- [1] N. Cabbibo, Phys.Rev.Lett. **10**, 531 (1963),
M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [2] A.B. Carter and A.I. Sanda, Phys. Rev. **D23**, 1567 (1981),
I.I. Bigi and A.I. Sanda, Nucl. Phys. **B193**, 85 (1981).
- [3] K. Abe *et al.* [Belle Collaboration], Phys.Rev.**D66**, 071102 (2002).
- [4] B. Aubert *et al.* [BABAR Collaboration], Phys.Rev.Lett. **89**, 201802 (2001).
- [5] T. Tomura, hep-ex/0305036, Presented at 38th Rencontres de Moriond on Electroweak Interactions and Unified Theories, Les Arcs, France, 15-22 Mar 2003.
- [6] M. Gronau and D. London, Phys.Rev.Lett. **65**,3381 (1990),
M. Gronau, D. London, N. Sinha and R. Sinha, Phys. Lett. **B514**, 315 (2001).
- [7] M. Gronau, O.F. Hernandez, D. London, J.L. Rosner, Phys.Rev. **D50**, 4529 (1994),
M. Gronau, J.L. Rosner and D. London, Phys.Rev.Lett. **73**, 21 (1994).
O.F. Hernandez, D. London, M. Gronau and J.L. Rosner, Phys. Lett. **B333**, 500 (1994).
- [8] M. Gronau, O.F. Hernandez, D. London, J.L. Rosner, Phys.Rev. **D52**, 6374 (1995).
- [9] M. Gronau and J.L. Rosner, Phys. Rev. Lett. **76**, 1200 (1996),
A.S. Dighe, M. Gronau and J.L. Rosner, Phys. Rev. **D54**, 3309 (1996),
M. Gronau and J.L. Rosner, Phys. Rev. **D57**, 6843 (1998).
- [10] C.S. Kim, D. London and T. Yoshikawa, Phys. Rev. **D57**, 4010 (1998).

- [11] R. Fleischer, Phys. Lett. **B365**, 399 (1996),
R. Fleischer and T. Mannel, Phys.Rev. **D57**, 2752 (1998),
R. Fleischer, Eur.Phys.J. **C6**, 451 (1999),
A.J. Buras and R. Fleischer, Eur.Phys.J. **C16**, 97 (2000).
- [12] M. Neubert, JHEP **9902**, 014 (1999).
- [13] R. Fleischer and J. Matias, Phys.Rev. **D66**, 054009 (2002).
- [14] M. Beneke, G. Buchalla, M. Neubert and C.T. Sachrajda, Nucl. Phys. **B606**, 245 (2001).
- [15] Y.-Y. Keum, H.-n. Li and A.I. Sanda, Phys. Lett. **B504**, 6 (2001),
Y.-Y. Keum, H.-n. Li and A.I. Sanda, Phys. Rev. **D63**, 054008 (2001).
- [16] C.-D. Lu and Z. Xiao, Phys.Rev. **D66**, 074011 (2002),
Z. Xiao, C.-D. Lu and L. Guo, hep-ph/0303070.
- [17] M. Gronau and J.L. Rosner, Phys.Rev. **D65**, 093012 (2002),
M. Gronau and J.L. Rosner, Phys.Rev. **D65**, 013004 (2002).
- [18] M. Gronau and J.L. Rosner, Phys.Rev. **D66**, 053003 (2002), Erratum-ibid. **D66**, 119901 (2002).
- [19] T. Yoshikawa, KEK-TH-878, to be published in Phys.Rev. **D** , hep-ph/0304038 .
- [20] Particle Data Group, K. Hagiwara *et al.*, Phys.Rev. **D66**, 010001-1 (2002).
- [21] A. Bornheim *et al.*, [CLEO Collaboration], hep-ex/0302026.
- [22] K. Abe *et al.* [Belle Collaboration], Phys.Rev. **D66**, 092002 (2002).
- [23] P.C. Bloom [BABAR Collaboration], eConf **C020805**, TTH06 (2002), hep-ex/0302030.
- [24] B. Aubert *et al.* [BABAR Collaboration], hep-ex/0206053,
B. Aubert *et al.* [BABAR Collaboration], hep-ex/0207065,
B. Aubert *et al.* [BABAR Collaboration], hep-ex/0303028,
B. Aubert *et al.* [BABAR Collaboration], Phys.Rev.Lett. **89**, 281802 (2002).
- [25] Y. Unno *et al.* [Belle Collaboration], hep-ex/0304035.
- [26] Y. Nir and H.R. Quinn, Phys.Rev.Lett. **67**, 541 (1991).
- [27] A.S. Dighe, T. Hurth, C.S. Kim and T. Yoshikawa, Nucl.Phys. **B624**, 377 (2002).